

Begonia semperflorens F? Sprint Plus

Maxi Mix

Item no.: C1274P

Crop Time

Spring: 9 - 11 weeks

Height

23cm

Exposure

Sun - Shade

Seed Form

Pelleted Seed

Best Uses

Bedding, Landscape

Culture guide

Usage

Packs, Pots, Hanging Baskets, Mixed Containers and Landscape

Sowing method

1-2 seeds or pellets per plug

Germination

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 6-8 days.

Temperature

Plug Culture:

72–76 °F (22–24 °C) days 1–11. For irrigation use warm water (above 64 °F / 18 °C) only.

Growing on:

68–70 °F (20–21°C) nights, 64–67 °F (18–19 °C) days for the first 14 days or until the roots reach the bottom of the container. Thereafter temperatures may be lowered to 62–65 °F (16–18 °C) day and night. An ADT (average daily temperature of 67 °F (19 °C) will give the fastest finished crop.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.