

Begonia tuberhybrida F₁

Nonstop Joy®

The Semi-Trailing & Multi-Purpose Begonia



Nonstop® Joy – The only Semi-Trailing Tuberous Begonia from Seed

- A multi-purpose begonia suited for premium patio pots and baskets
- Extra large double flowers
- Well-branched plants
- Blooms all summer long
- Easy to produce and easy to ship

Technical Guide: [Click here](#)

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Spring: 16 - 18 weeks
Height ∅	9 " / 23 cm
Width ∅	9 " / 23 cm
Exposure	Partial shade - Shade
Seed Form	Pelleted Seed
Best Uses	Bedding, Hanging basket, Pot Plant

CULTURE GUIDE

Begonia tuberhybrida F₁ Nonstop Joy®

Usage

Pots, Hanging Baskets, Mixed Containers and Landscape

Sowing method

Sow 1-2 seeds or pellets per plug.

Germination

Maintain optimal conditions for seedling development, should begin on the day of sowing until root emergence. Expect root emergence in 7-10 days.

Media

Plug Culture: pH 5.5-5.8; EC 0.5. Begonias are sensitive to high salt levels. Use a well-drained, growing substrate. pH 5.5-5.8; EC 1.25-1.5.

Growing on: Use a well-drained, growing substrate. pH 5.5-5.8; EC 1.0-1.5.

Temperature

Plug Culture: Maintain 22-23°C (72-73°F) until root emergence. Higher temperatures, exceeding 27 °C (80°F) will inhibit germination. Upon radicle emergence, on day 10-14 reduce the temperature to 20-21°C (68-70°F) until cotyledon expansion. On day 21 the temperature can be reduced further to 20°C (68°F). For irrigation use warm water (above 18°C/ 64 °F) only. Maintain the ideal temperature to achieve the shortest crop time on days 28-42 keeping the temperature 20-21°C (68-70°F). A slightly lower temperature of 19-20°C (66-68°F) will reduce stretch in the seedlings. Tuberous begonias are very responsive to DIF where a 1-2 °C (34-26°F) DIF will also keep plants compact. Once roots have reached the bottom of the tray, (after day 42), the temperature can be lowered to 18-19°C (64-66°F).

Growing on: After transplanting maintain 18-20°C (64-68°F) nights for the first 14 days or until the roots reach the bottom of the container. Thereafter temperatures may be lowered to 16-18°C (60-64°F). An ADT (average daily temperature of 19°C (66°F) will give the fastest finished crop. Temperatures below 14°C (57°F) will result in tuber formation and delay the crop. A DIF of 1-2°C (34-36°F) will result in a more compact crop requiring little to no growth regulators.

Fertilization

Begin fertilizing early once germination is complete, approximately day 14 lower rates of feeding at 50 ppm 2-3 times per week will help to size up the seedlings. Under higher light conditions use a 20-10-20 fertilizer and under lower light a 17-5-17. Once seedlings are established the 17-5-17 fertilizer works well. Include a micro nutrient package to give adequate supply of minor elements. Alternate between a calcium-based fertilizer (14-4-14 or 15-5-15) and an ammonium-based fertilizer (17-5-17) at 50- 100 ppm nitrogen every 2-3 waterings. Fertilizer applications can be gradually increased in the later stages of bulking where a constant feed at 100 ppm is used. Under high light and long days an ammonium-based fertilizer 20-10-20 at 50-100 ppm can also be used.

Growing on: Alternate between calcium-based fertilizer 14-4-14 and an ammonium fertilizer 17-5-17 at 100-150 ppm. Keep the media EC at 1.5. Application of potassium nitrate can help to keep the plants more compact. Under higher light and warmer temperatures a fertilizer with additional ammonium can be used. Tall, stretched plants with few flowers indicate too much ammonium. Stunted, chlorotic plants with marginal leaf burn indicate a lack of calcium and magnesium. Under high light and extended day length an ammonium-based feed (20-10-20) at 100-150 ppm nitrogen can also be used.

Tip

Nonstop Joy® has excellent branching and continuous flowering all season long. Will actually finish 5-7 days earlier than the other Nonstop® series.

COLORS OF THE SERIES

Begonia tuberhybrida F₁ Nonstop Joy[®]



Mocca Rose
BT0609P



Mocca White
BT0606P



Orange
BT0601P



Peaches & Dreams
BT0604P



Red
BT0602P



Rose Picotee
BT0603P



Yellow
BT0607P



Mix
BT0699P