




Impatiens walleriana F<sub>1</sub>

# Lollipop

### **Fervent Fruit Mix**

Item no.: IW0195R



#### The Perfect POP Series for Color

- Uniform series in flowering and habit
- Full plants hide the pot
- Excellent branching more flowers!
- Intense colors "pop" at retail
- No stretching holds well

#### **Technical Guide: Click here**

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Only available in North America.

Crop Time	Spring: 8 - 11 weeks
Height ∅	13 " / 33 cm
Width Ø	11 " / 28 cm
Exposure	Partial shade
Seed Form	Raw Seed
Best Uses	Bedding, Landscape



# **CULTURE GUIDE**

Impatiens walleriana F₁ Lollipop

#### Usage

Packs, Pots, Hanging Baskets, Mixed Containers and Landscape

#### Sowing method

1 seed per plug

#### Germination

Optimum conditions for seedling development, beginning on the day of sowing until radical emergence Expect radicle (root) emergence in 3-5 days.

#### **Temperature**

72-75  $^{\circ}$ F (22-24  $^{\circ}$ C) until day 6 then reduce slightly to 70-72  $^{\circ}$ F (21-22  $^{\circ}$ C). A germination temperature below 70  $^{\circ}$ F (21  $^{\circ}$ C) will slow the speed and uniformity of germination. Excessive temperature above 77  $^{\circ}$ F (25  $^{\circ}$ C) can cause a thermodormancy. Temperatures below 65  $^{\circ}$ F (18  $^{\circ}$ C) may cause tip abortion and malformed seedlings.

#### **Fertilization**

Begin fertilizing early on day 3 with low rates of 25-50 ppm nitrogen using a calcium based feed (14-0-14). Maintain an EC less than 0.75 for the first 7-10 days.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons. Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.



# **COLORS OF THE SERIES**

## Impatiens walleriana F<sub>1</sub> Lollipop



Bubblegum (Pink) IW0104R



Cherry (Red) IW0105R



Coconut (White)
IW0111R



**Dragon Fruit (Lipstick)**IW0107R



Fruit Punch (Rose) IW0106R



Orange IW0103R



Peach (Salmon) IW0108R



Pink Lemonade (Light Pink) IW0102R



Pomegranate (Carmine) IW0101R



Raspberry (Violet)
IW0110R



Tangerine (Deep Salmon) IW0109R



Fervent Fruit Mix IW0195R



Maxi Mix IW0199R