



Petunia x hybrida F₁

SUCCESS![®] 360°

Purple

Item no.: PH0507R/P



Impressive Show Maker from 360°

- Round growing grandiflora Petunia
- Very uniform series in plant habit
- Early flowering with large, attractive flowers
- Wide range of colors with Star shades
- Fast filling of packs, pots or baskets

Technical Guide: [Click here](#)

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Spring: 9 - 13 weeks
Height ∅	13 " / 33 cm
Width ∅	11 " / 28 cm
Exposure	Sun
Seed Form	Raw Seed, Pelleted Seed
Best Uses	Hanging basket, Landscape

CULTURE GUIDE

Petunia x hybrida F₁ SUCCESS!® 360°

Usage

Packs, Pots, Hanging baskets, Mixed Containers and landscape

Sowing method

1 pellet per plug

Germination

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 3-5 days.

Media

Begin by watering to saturated (5); applying enough water to help dissolve the pellets. After sowing do not allow the pellets to dry back before moving to the germination chamber or benches. Maintain saturation (5) for 3-4 days or until radicle emergence. On day 5 reduce media moisture to wet (4) for the next 5-6 days. On day 10 reduce the moisture further to medium (2). Alternate between wet (4) and a medium (2) between watering.

Temperature

Plug Culture: 72-76 °F (22-24 °C) until radicle emergence and then reduce to 68-70 °F (20-21 °C). The temperature can be lowered on approximately day 5. Once cotyledons have fully expanded lower the temperature further to 65-68 °F (18-20 °C). Growing On: After transplant, maintain temperatures > 55 °F (13 °C) nights for the first 3-4 weeks to initiate flower bud development. The night temperatures can be lowered further to 50 °F (10 °C) to encourage basal branching and compactness. However, lower temperatures may also substantially decrease the number of flowers initiated. Growing at cooler temperatures will produce a higher quality plant. An ADT (average daily temperature of 67 °F (19 °C) will give the fastest finished crop.

Fertilization

Maintain an EC < 1.0. Fertilized water should not exceed an EC of 0.5. Upon initial germination, approximately day 5-6 begin feeding with 50 ppm nitrogen. Pay attention to the addition of boron since low boron can cause tip abortion. Ideal boron concentration is 0.5 ppm.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

COLORS OF THE SERIES

Petunia x hybrida F₁ SUCCESS!® 360°



Blue
PH0506R/P



Burgundy
PH0508R/P



Burgundy Vein
PH0510R/P



Deep Pink
PH0513R/P



Deep Rose
PH0504R/P



Deep Salmon
PH0502R/P



Light Yellow
PH0515R/P



Pink
PH0503R/P



Purple
PH0507R/P



Purple Vein
PH0509R/P



Red
PH0505R/P



Red Star
PH0512R/P



Rose Star
PH0511R/P



Violet
PH0514R/P



White
PH0501R/P



Maxi Mix
PH0599R/P