

#### Portulaca grandiflora F<sub>1</sub>



### Orange

Item no.: PG2205R/U



- Short crop time, fast growing
- Vigorous, well branched habit
- For pots, packs and baskets
- Wide color range with exciting tones

#### Technical Guide: Click here

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Spring: 10 - 12 weeks
Height Ø	5 ″ / 13 cm
Width Ø	11 ″ / 28 cm
Exposure	Sun
Seed Form	Raw Seed, Multipelleted Seed
Best Uses	Bedding, Landscape, Rockery



# **CULTURE GUIDE**

Portulaca grandiflora F1 Sundial

#### Usage

Mixed containers, bedding, landscaping

#### Sowing method

2 seeds per plug, 1 multi-seed pellet per plug

#### Germination

Stage I: 2-3 days at 70-78 °F (22-26 °C). Sow in media with very low soluble salt levels and pH 5.5-6.2. Light is not essential, but beneficial for germination. Keep soil moist but not wet. Keep relative humidity between 90-95%. At the end of stage I the radicle will have emerged and the cotyledons begin to unfurl.

#### Growing on

4-5 weeks after sowing (288-tray) transplant 1 plant into 10-11 cm (4-5") pots or packs

#### Media

Use a well-drained substrate with pH 5.5 -6.2 and 1-1.5kg/m<sup>3</sup> fertilizer

#### Temperature

18 °C-22 °C (65-72 °F). Temperatures below 18 °C (65 °F) will inhibit induction and result in poor or no vegetative growth.

#### Fertilization

Moderate fertilization levels are required. Fertilize the crop with 150-200 ppm nitrogen, using a complete balanced fertilizer.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons. Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair. Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable. Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.



## **COLORS OF THE SERIES**

### Portulaca grandiflora F<sub>1</sub> Sundial



Chiffon PG2201R/U



Coral



**Light Pink** PG2208R/U



White PG2213R/U





Orange PG2205R/U



Yellow PG2214R/U



Cream PG2202R/U



Peppermint PG2206R/U



**Caliente Mix** PG2298R/U



**Deep Red** PG2212R/U



Pink PG2207R/U



Mix PG2299R/U



Fuchsia PG2203R/U



Scarlet PG2211R/U



**Peach Mix** PG2295R/U



Gold PG2204R/U



Tangerine PG2209R/U



**Valentine Mix** PG2297R/U



**Volcano Mix** PG2296R/U

