



1. [Home](#)

Viola wittrockiana F?

## Inspire® Plus

### Most Advanced Breeding in Large-Flowered Pansies

- Ideal for spring and fall production
- Uniform flower timing and habit across the series
- Short flower stems, compact plants – no stretching!
- Outstanding seedling quality
- BeGreen Primed – chemical free and eco-friendly

**Technical Guide:** [Click here](#)

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.



[Bookmark](#)

[Recommend](#)

[Print](#)

### **Crop Time**

Spring: 25 - 26 weeks , Autumn: 10 - 12 weeks

### **Height ?**

7 ? / 18 cm

### **Width ?**

7 ? / 18 cm

### **Exposure**

Sun

### **Seed Form**

Raw Seed, BeGreen Priming

### **Best Uses**

Bedding, Landscape

## Culture guide

### Usage

Packs, Pots, Mixed Containers and Landscape/Mass plantings

### Sowing method

1 seed per plug

### Germination

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 2-4 days.

### Temperature

65-68 °F (18-20 °C) until radicle emergence, then lower the temperature gradually to 62-65 °F (17-18 °C).

### Fertilization

Begin feeding early using a calcium based fertilizer at lower rates to keep the an adequate amount of calcium and nitrogen supplied to the seedlings. On days 5-7 begin feeding with a calcium based fertilizer (14-2-14, 13-2-13, 15-5-15, 17-5-17) at 50-60 ppm. Maintain the EC between 0.5 and 0.75. Keep phosphorous levels between 6-8 ppm and boron supplied at 0.5 ppm.

---

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the

young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

[Download](#)