

Delphinium hybrida

Benary's Pacific

Guinevere

Item no.: DH0108R



- FastraX perennial: First year flowering without vernalization
- Rigorously tested for excellent flower and stem quality
- Excellent cut flower with long, impressive flower stems
- Long vase life

Crop Time	Spring: 18 - 22 weeks
Height ∅	63 " / 160 cm
Exposure	Sun
Seed Form	Raw Seed
Heat Zone	6-1
Hardiness Zone	3b-7b
Best Uses	Bedding, Cutflower, Landscape

CULTURE GUIDE

Delphinium hybrida Benary's Pacific

Usage

Background plant, cut flower

Sow time

Outdoor forcing: March-April; Indoor forcing: December-May

Sowing method

2-3 seeds per plug

Germination

20-30 days at temperatures of 59-68 °F (15-18 °C).

Growing on

Grow on at 60-65 °F (15-18 °C)

Media

Use a well-drained, growing substrate with 10-30 % clay, 0-15 % parts (e.g. bark, sand, perlite), 1-2 kg /m³, complete balanced fertilizer. 2-4 kg/m³ slow release fertilizer (3-9 months), iron-chelate, micronutrients, pH 5.5-6.5. Field: well-drained, moderate nutrition supply, humus, loamy soils. Standard fertilization: 60-80 g/m² slow release fertilizer (potassium balanced). Do not improve the soil with peat. Keep the soil open.

Temperature

Grow at 15-18 °C (59-64,5 °F) or outdoors. In winter indoors frost free at 3-5 °C (37,5-41 °F) or outdoors. Outdoor fleece cover and dry substrate needed. In spring the plants start to grow for 10-12 weeks at 15-18 °C (59-64,5 °F). A chilling period for flower initiation is not required.

Fertilization

High fertilization levels are required. Fertilize the crop weekly with 100-150 ppm nitrogen (at 2 kg/m³ slow release fertilizer in substrate), using a complete balanced fertilizer. Avoid high ammonium and high nitrogen levels. Don't fertilize after flowering or after early September. In spring fertilize with 100-150 ppm nitrogen using a complete balanced fertilizer. Prevent magnesium deficiency by applying magnesium sulphate (0,05 %) 1-2 times and in case of Iron deficiency apply iron-chelate for 1-2 times. The fertilization with slow release fertilizer is preferred to fertilization with complete balanced fertilizer, because the roots are very sensitive to high salt levels in substrates. Avoid high fertilizer concentrations, it is advisable to fertilize several times with low concentrations weekly. Field: After harvest fertilize the plant additionally with 60-80 g/m² of a slow release and potassium balanced fertilizer. Don't fertilize after flowering.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

COLORS OF THE SERIES

Delphinium hybrida Benary's Pacific



Astolat
DH0109R



Black Knight
DH0101R



Blue Bird
DH0106R



Blue Jay
DH0107R



Cameliard
DH0102R



Galahad
DH0110R



Guinevere
DH0108R



King Arthur
DH0104R



Percival
DH0105R



Summer Skies
DH0103R



Mix
DH0199R