

Delphinium hybrida Benary's Pacific Blue

Bird

Item no.: K2831

Crop Time

Spring: 18 - 22 weeks

Height

160cm

Exposure

Sun

Seed Form

Raw Seed

Hardiness Zone

3b-7b

Best Uses

Bedding, Cutflower, Landscape

Culture guide

Usage

Background plant, cut flower

Sow time

Outdoor forcing: March-April; Indoor forcing: December-May

Sowing method

2-3 seeds per plug

Germination

20-30 days at temperatures of 59-68 °F (15-18 °C).

Growing on

Grow on at 60-65 °F (15-18 °C)

Media

Use a well-drained, growing substrate with 10-30 % clay, 0-15 % parts (e.g. bark, sand, perlite), 1-2 kg /m³, complete balanced fertilizer. 2-4 kg/m³ slow release fertilizer (3-9 months), iron-chelate, micronutrients, pH 5.5-6.5. Field: well-drained, moderate nutrition supply, humus, loamy soils. Standard fertilization:60-80 g/m² slow release fertilizer (potassium balanced). Do not improve the soil with peat. Keep the soil open.

Temperature

Grow at 15-18 °C (59-64,5 °F) or outdoors. In winter indoors frost free at 3-5 °C (37,5-41 °F) or outdoors. Outdoor fleece cover and dry substrate needed. In spring the plants start to grow for 10-12 weeks at 15-18 °C (59-64,5 °F). A chilling period for flower initiation is not required.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.