

Pentas lanceolata F₁

Graffiti[®]

Lipstick

Item no.: PL0417P



The Unbeatable Series on the Market!

- Profit from the widest range of colors in a Pentas series:
17 intense colors & 2 mixes to choose from!
- Uniformity in height and flower timing
- Well-branched plants with dense flowers for maximum color at retail
- Excellent heat tolerance and garden performance
- First bicolor tones in seed Pentas

Technical Guide: [Click here](#)

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Spring: 13 - 16 weeks
Height ∅	12 " / 30 cm
Width ∅	11 " / 28 cm
Exposure	Sun
Seed Form	Pelleted Seed
Best Uses	Bedding, Landscape, Pot Plant

CULTURE GUIDE

Pentas lanceolata F₁ Graffiti®

Usage

Pots, Mixed Containers and Landscape

Sow time

January-April

Sowing method

1 pellet per plug. No cover is necessary.

Germination

Maintain optimal conditions for seedling development beginning on the day of sowing until radical emergence. Expect radical emergence in 7-10 days. Humidity should be between 95-100% until day 10; then reduce to 40-60%.

Growing on

Day neutral plant, will flower regardless of day length. Very responsive to irradiance and additional lighting. Providing a 14-16 hrs. day length, especially in the seedling stages, will shorten the crop significantly. In addition, growing at warmer temperatures will shorten the crop time.

Media

Plug culture: pH 6.2-6.5 Starting with the proper pH of the media will improve the performance of the seedlings. Pentas can exhibit iron toxicity at lower pH levels, below 5.5. Pentas require close attention to the proper media pH. If the pH is too high, a micro nutrient deficiency may occur and if too low, an iron toxicity can occur. EC < 0.5.

Temperature

Plug culture: 23-26°C (74-78°F). Once germination is completed with fully expanded cotyledons, on day 14 the temperature can be lowered slightly to 22°C (72°F). Water trays using tempered water with a minimum temperature of 18°C (64°F).

Media temperatures below 16°C (60°F) will inhibit the germination and growth. 20-21°C (68-70°F). As plants become more mature the temperature can be lowered to 18-20°C (64-68°F) nights and 22-23°C (72-73°F) days. Warmer temperatures will benefit and shorten the finish time.

Growing on: 20-21°C (68-70°F) nights, 22-23°C (72-73°F) days for the first 14 days or until the roots reach the bottom of the container. Thereafter temperatures may be lowered to 16-18°C (60-64°F) nights and 20-23°C (68-74°F) days. Higher temperatures are beneficial and will shorten the crop time. Pentas do not seem to have a maximum temperature that will inhibit growth and flowering.

Fertilization

Plug culture: Maintain an EC < 0.75. At this stage fertilized water should not exceed an EC of 0.5 Begin feeding on day 10 with 50 ppm 14-2-14, 14-4-14 or 17-5-17. Keep phosphorous levels < 8 ppm, iron levels at 2-3 ppm. Maintain the EC levels below 1.2. Under lower light conditions fertilize with a calcium based fertilizer, 14-4-14 at 100 ppm. Under higher light use a 17-5-17 feed at 100 ppm.

Growing on: : Under low light conditions fertilize with a 14-4-14 fertilizer at 100-150 ppm and under high light conditions use a 17-5-17 fertilizer at 100-150 ppm. Watch for calcium and magnesium deficiencies which can cause stunted plants.

Tip

Graffiti® is the earliest, most uniform and compact Pentas series on the market. It has the broadest color range and garden performance that lives up to the Graffiti® brand name growers and landscapers trust.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable

COLORS OF THE SERIES

Pentas lanceolata F₁ Graffiti®



Appleblossom
PL0402P



Bright Red
PL0419P



Cranberry
PL0405P



Deep Red
PL0413P



Flirty Pink
PL0403P



Fuchsia
PL0409P



Lavender Pink
PL0412P



Lipstick
PL0417P



Neon Plum
PL0415P



Pink
PL0418P



Red Velvet
PL0422P



Rose
PL0420P



Ruby
PL0406P



True Pink
PL0407P



Ultra Violet
PL0410P



Violet
PL0421P



White
PL0411P



Maxi Mix
PL0499P



Selected Mix
PL0498P