

Begonia tuberhybrida F₁

Nonstop[®] Mocca

Deep Orange

Item no.: BT0503P



Unique - the Only Dark Leaved Tuberous Begonia Series on the Market

- Full, compact and rounded plant habit
- Uniform and strong branching plants
- Excellent outdoor performance
- Highest transplantable seedlings in the industry
- Free flowering of large double flowers
- Comprehensive color assortment

Technical Guide: [Click here](#)

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Spring: 15 - 18 weeks
Height ∅	10 " / 25 cm
Width ∅	9 " / 23 cm
Exposure	Partial shade - Shade
Seed Form	Pelleted Seed
Best Uses	Bedding, Landscape, Pot Plant

CULTURE GUIDE

Begonia tuberhybrida F₁ Nonstop® Mocca

Usage

Bedding, patio containers and landscape, window boxes, pot plants

Sow time

November for flowering pots from April onwards, December-January for flowering bedding plants from May onwards

Sowing method

1-2 seeds per plug

Germination

Germination will occur in 7-14 days at 75-78 °F (23-25 °C). Sow seed on a fine media with good water holding capacity and good drainage. Consistent moisture levels are important to uniform germination. Humidity levels above 95 % and a media pH between 5.5 and 6.5 are important. Do not cover seed as light is required to germinate. Supplemental 24-hour assimilation light provided at this stage will increase germination, reduce crop time and improve plug quality.

Growing on

Transplant plugs into finished containers with a well drained media, and pH of 5.5 to 6.5. Maintain day length in excess of 14 hours. Continued supplemental lighting will improve plant quality and shorten crop time. Growing temperatures between 68-72 °F (18-22 °C) optimize growth and flowering. Fertilize at 150-250 ppm nitrogen in a well-balanced formula.

Media

Use a well-drained, growing substrate, pH: 5.5-6.2.

Temperature

Grow at 16-18 °C. 10 days before selling the temperatures can be decreased to 16 °C. Temperatures below 14 °C will result in tuber formation and crop delay.

Fertilization

Moderate fertilization levels are required. At high nitrogen levels the foliage can become very big. Avoid pH above 6.0, as high pH causes iron deficiency. Apply chelated iron, if chlorosis becomes a problem. Avoid high salt levels in substrate. It is advisable to fertilize several times with low concentrations weekly.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

COLORS OF THE SERIES

Begonia tuberhybrida F₁ Nonstop® Mocca



Bright Orange
BT0502P



Cherry
BT0510P



Deep Orange
BT0503P



Deep Red
BT0505P



Light Pink Shades
BT0513P



Pink Shades
BT0507P



Red
BT0511P



Scarlet
BT0504P



White
BT0506P



Yellow
BT0501P



Mix
BT0599P